

On the external photon fields in Fermi bright blazars

Xiaoyan Gao^{1,2,3}, Jiancheng Wang^{1,2} and Ming Zhou^{1,2,3}

¹ National Astronomical Observatories/Yunnan Observatories, Chinese Academy of Sciences, Kunming, Yunnan province 650011, P.R. China xy-gao@ynao.ac.cn

² Key Laboratory for the Structure and Evolution of Celestial Objects, Chinese Academy of Sciences, Kunming, Yunnan province 650011, P.R. China

³ Graduate School, Chinese Academy of Sciences, Beijing, P.R. China Received [year] [month] [day];

accepted [year] [month] [day]

Abstract The external Compton (EC) model is used to study the high energy emission of some blazars, in which the external photon field is considered to dominate inverse Compton radiation. We explore the properties of external photon field through analyzing the *FERMI* LAT bright AGN sample within three months detection. In the sample, assuming the high energy radiation of low synchrotron peaked blazars from the EC process, we find that the external photon parameter $U_{\text{ext}}/\nu_{\text{ext}}$ may not be a constant. Calculating synchrotron and inverse Compton (IC) luminosity from the quasi-simultaneous broadband spectrum energy distributions (SEDs), we find that they have an approximately linear relation. This indicates that the ratio of external photon and magnetic energy density is a constant in the comoving frame, implying that the Lorentz factor of the emitting blob depends on external photon field and magnetic field. The result gives a strong constraint on the jet dynamic model.

Key words: galaxies: active – galaxies: jets – gamma-rays: galaxies – radiation mechanism: non-thermal

1 INTRODUCTION

Blazars, presenting weak (or even absent) broad emission line and large optical polarization, are radio loud subclass of active galactic nuclei (AGNs). They are thought to be objects emitting non-thermal radiation across the entire electromagnetic spectrum from a relativistic jet which is closely along the line of sight and hence appears Doppler boosted (Begelman & Rees (1984); Urry (1999)). The broadband spectrum energy distribution of blazars are double bumped with a first peak usually in the soft-to-medium X-ray range, and a second one in GeV-TeV energies band (Sambruna et al.(1996)). The first peak is generally dominated by synchrotron emission (Marscher & Gear (1998)), and the second one is uncertain.

The high energy emission is usually attributed by Compton upscattering of soft photons. The soft photons may come from synchrotron process called synchrotron self Compton (SSC) model (Maraschi et al.(1992); Marscher & Gear(1998)), or from external field photons which is called External Compton (EC) model. The external field photons can possibly include accretion disk radiation (e.g., Dermer et al.(1992); Dermer & Schlickeiser(1993)), broad line region (BLR) (Sikora et al.(1994); Blandford & Levinson(1995); Ghisellini & Madau (1996); Dermer et al. (1997)), dust torus (Blażejowski et al.(2000)) or synchrotron emission from other regions(Georganopoulos & Kazanas (2003); Ghisellini & Tavecchio(2008)). All these different scenarios have been tested by specific sources.

Blazars contains BL Lac objects and flat spectrum radio quasars (FSRQs). In previous literatures, BL Lac objects are often subdivided into two or three subclass depending on their SED which was first introduced by Padovani & Giommi (1995). Depending the peak energy of the synchrotron emission, BL Lacs are classified into low-frequency or high-frequency BL Lac objects, called LBL and HBL respectively. Abdo et. al. (Abdo et. al. 2010) give a new classification to all types of non-thermal dominated AGNs. They are low synchrotron peaked (LSP), intermediate synchrotron peaked (ISP) and high synchrotron peaked (HSP) blazars, defined by the location of the low energy SED peak. The LSP sources, consisting of flat spectrum radio quasars and low frequency peaked BL Lac objects (LBLs), have the synchrotron peak in the far-infrared or infrared regime with $\nu_{\text{peak}}^S \leq 10^{14}$ Hz. The ISP sources, mainly including LBLs and intermediate BL Lac objects (IBLs), have their synchrotron peak in optical-UV frequencies with $10^{14} \leq \nu_{\text{peak}}^S \leq 10^{15}$ Hz, while the HSP objects, almost all known to be high-frequency-peaked BL Lac objects (HBLs), have their synchrotron peak at X-ray energies with $\nu_{\text{peak}}^S > 10^{15}$ Hz.

In the framework of leptonic models, the blazar sequence $\text{FSRQ} \rightarrow \text{LBL} \rightarrow \text{IBL} \rightarrow \text{HBL}$ is thought to be a decreasing contribution of external radiation fields to radiative cooling of electrons and production of high energy radiation (Ghisellini et al. (1998); Celotti & Ghisellini (2008)). The γ ray radiation of FSRQs always thought to be attributed by EC model (Cao & Bai, 2008). In this paper we use the sample of LSP blazars from the first three month observation of *Fermi Gamma Ray Space Telescope* to study the properties of high energy emission and external photon field based on the EC model. The peak frequencies and fluxes are obtained through fitting the SEDs which are quasi-simultaneous multifrequency measurement instead of using a empirical method from spectral slopes α_{ox} and α_{ro} . We find that the external photon parameter in the observer frame $U_{\text{ext}}/\nu_{\text{ext}}$ is not constant. We also find that there is a linear correlation between synchrotron luminosity L_S and inverse Compton (IC) luminosity L_{IC} , implying that the ratio between the external radiation and the magnetic field energy densities inside the source (in the comoving frame) depends on Lorentz factor. In section 2 we present the sample of LSP sources. In section 3 we show the statistical method and results. The discussion the results are shown in section 4.

We adopt a concordance cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.7$ and $\Omega_M = 0.3$.

2 THE SAMPLE

The Large Area Telescope (LAT) on board the *Fermi Gamma Ray Space Telescope*, launched on 2008 June 11, provides unprecedented sensitivity in γ -ray band (20 MeV to over 300 GeV, Atwood et. al. (Atwood et. al. 2009)). The first three months, from 2008 August 4 to October 31, of operations in the sky-survey mode led to the compilation of a list of 106 high confidence γ -ray sources. Abdo et. al. (Abdo et. al. 2010) give the quasi-simultaneous SEDs of 48 blazars from the LAT observation. They include 23 FSRQs, 19 BL Lacs and one unknown type blazar, which have certain redshifts.

The sample we adopted consists of 30 low synchrotron peak (LSP) blazars, including 23 FSRQs and 7 low energy peak BL Lacs. In view of blazar sequence, the gamma-ray emission of these LSPs could be dominated by external compton radiation. We also present 8 high energy peak BL Lacs for contrast. Usually the HBLs lack the emission line, the SSC radiation could dominate their gamma-rays. All sources are listed in table 1. Column 1 gives the source name presented by Abdo et al. (Abdo et. al. 2009); Column 2 gives the redshift; Column 3, 4, 5 and 6 are the peak frequencies and fluxes of synchrotron and inverse Compton (IC) emission. We use the peak frequencies ν_{peak}^S , $\nu_{\text{peak}}^{\text{EC}}$, and the peak fluxes $\nu_{\text{peak}}^S F(\nu_{\text{peak}}^S)$, $\nu_{\text{peak}}^{\text{EC}} F(\nu_{\text{peak}}^{\text{EC}})$, given by Abdo et. al. (Abdo et. al. 2010) who using a simple third-degree polynomial to fit the SED (Kubo et al. (1998)), where the data are quasi-simultaneous multifrequency observation. Column 7 lists the total synchrotron luminosity; Column 8 is the total IC luminosity; Column 9 is the IC luminosity calculated by the EC model; Column 10 gives the classifications.

3 MODEL AND TEST

We consider the homogeneous EC model to study the properties of high energy SED for LSP sources. The synchrotron and IC peak frequencies are given by (Tavecchio et. al.(1998))

$$\nu_{\text{peak}}^S = \frac{4}{3} \nu_L \gamma_b^2 \delta, \quad (1)$$

and

$$\nu_{\text{peak}}^{\text{EC}} = \frac{4}{3} \nu_{\text{ext}} \gamma_b^2 \Gamma \delta, \quad (2)$$

where $\nu_L = eB/(2\pi m_e c)$ is the Larmor frequency, γ_b is the Lorentz factor of electron spectral break, and ν_{ext} is the concentrated frequency of external photons. Γ is the Lorentz factor of the emitting blob. While δ is the Doppler factor defined by $\delta = 1/(1 - v/c \cos \theta)$, v is the relativistic bulk velocity, and θ is the angle of the jet relative to observer's line of sight.

The ratio between the total luminosity of synchrotron and IC radiation is related to the ratio between external field and magnetic energy density as

$$\frac{L_{\text{IC}}}{L_S} = \frac{U'_{\text{ext}}}{U'_B} \simeq \frac{17}{12} \frac{\Gamma^2 U_{\text{ext}}}{U'_B}, \quad (3)$$

where $U'_{\text{ext}} \simeq (17/12) \Gamma^2 U_{\text{ext}}$ is the external photon energy density in the comoving frame (Ghisellini & Madau (1996)), and U'_B is the magnetic field energy densities inside the source (in the comoving frame). Here we declare that parameters with prime (" ' ") presents in coming frame, otherwise are in observer frame.

From the above equations we obtain

$$\frac{L_{\text{IC}}}{L_S} \simeq \frac{17e^2}{6\pi m_e^2 c^2} \frac{U_{\text{ext}}}{\nu_{\text{ext}}^2} \left(\frac{\nu_{\text{peak}}^{\text{EC}}}{\nu_{\text{peak}}^S} \right)^2. \quad (4)$$

The SEDs of the sources in the sample are fitted by Abdo et al. (Abdo et. al. 2010). We use the SEDs to obtain the differential luminosity of synchrotron and IC radiation. We then calculate the total luminosity L_S and L_{IC} by integrating the differential luminosity. The results are shown in column (7) and column (8) of table 1. Assuming L_{IC} from external Compton emission mainly, we examine the relation of L_{IC}/L_S and $\nu_{\text{peak}}^{\text{EC}}/\nu_{\text{peak}}^S$ shown in Figure 1, and find that the relation is rather disperse. No obvious correlation implies that the parameter $U_{\text{ext}}/\nu_{\text{ext}}^2$ of the external photons is not a constant from equation (4). The result is different with common scenario on the external photon from the BLR (Ghisellini et al. (1998); Celotti & Ghisellini (2008); Ghisellini & Madau (1996); Tavecchio & Ghisellini(2008)). In particular, we take the common values of $U_{\text{BLR}} = 2.65 \times 10^{-2} \text{ erg cm}^{-3}$ and $\nu_{\text{ext}} = 2 \times 10^{15} \Gamma \text{ Hz}$, and calculate the EC luminosity L_{EC} from L_S using the equation (4) shown in column (9). We find that the L_{EC} of many sources is larger than the L_{IC} from the SEDs, indicating that the previous results on the external photon field have questionable validity.

We also show the relation between L_{IC} and L_S in Figure 2, indicating a significant correlation. The best fit to the $L_{\text{IC}} - L_S$ relationship is

$$\text{Log} L_{\text{IC}} = 1.1 \text{Log} L_S - 4.5 \quad (5)$$

with a correlation coefficient $R = 0.58$, and the relationship is nearly linear, spanning a wider range of both luminosity. The relation indicates that L_{IC}/L_S is an approximate constant, implying the ratio of external photon and magnetic energy density to be constant in the comoving frame. If the gamma-ray blob is inside the BRL, from equation (3), we will obtain $\Gamma \propto (U'_B/U'_{\text{ext}})^{1/2}$, implying that the motion of the jet could relate to the magnetic field and external photon. Usually the magnetic field takes an important role in the acceleration and collimation of the relativistic jet (Leismann et al. (2005), McKinney




Fig. 1 The relation between L_{IC}/L_S and $\nu_{\text{peak}}^{\text{EC}}/\nu_{\text{peak}}^{\text{S}}$. Triangles: FSRQs (LSPs); black circles: BL Lac objects (LSPs) and red circles: BL Lacs which are HSPs. The dashed line is the fitted line with the slope of 2.

(2006), Keppens et al. (2008)), while the external photons can affect the jet dynamics through radiative drag (Sikora et. al. (1996), Luo & Protheroe (1999)). Therefore, the relation we obtained provides an observational evidence for the balance between the acceleration and deceleration of the jet. The main contribution for high energy radiation of HSPs is always considered to be the SSC emission, and the EC emission could be a supplement for high energy radiation. For comparison, we plot HSPs in Figure 2 and find that HSPs behave similarly with LSPs, locating at low luminosity range.

4 CONCLUSION AND DISCUSSION

The BL Lac objects in our sample include two subsamples: LBLs and HBLs. The LBLs together with FSRQs is considered as LSPs (Abdo et. al.(Abdo et. al. 2010)) as mentioned above. External Compton scattering is likely to be the main emission process responsible for the copious amounts of γ -rays produced in LSP blazars. The soft seed photons outside the γ -ray emitting region dominate gamma-ray emission. Usually the HBLs lack the broad emission line, the SSC could dominate their gamma-rays. The equation (3) is obviously invalid for the HBLs, while the HSPs in the figure 1 only show their position for comparison with other sources and don't present their relations with gamma-ray emission mechanism. For LBLs, if the gamma-rays originate from the blob inside the BLR and the size of the BLR is proportional to $L^{0.5}$ (Celotti & Ghisellini(2008)), the energy density and peak frequency of external photons will be a constant with good approximation. It is indicated that the ratio between EC and synchrotron luminosity just depends on the peak frequency $(\nu_{\text{peak}}^{\text{S}}/\nu_{\text{peak}}^{\text{EC}})^2$. In the paper we have




Fig. 2 The relation of the total luminosity L_S and L_{IC} . Symbols are the same as in figure 1.

presented the statistic analysis of a sample of 30 LSP blazars from the *Fermi*-LAT observation. After examining the relation between L_{IC}/L_S and $\nu_{\text{peak}}^{\text{EC}}/\nu_{\text{peak}}^{\text{S}}$ for LSP blazars, we find there is almost no correlation between them. It means that the parameter $U_{\text{ext}}/\nu_{\text{ext}}^2$ of the external photons is not a constant or the equation (3) is invalid for some LSPs due to the gamma-ray blob locating outside the BLR. Our statistical results show that the gamma-rays of LSPs could be produced at different external photon fields. There are two kinds of emissive scenarios. (1) The gamma-ray blob is inside the BLR, the figure 1 implies that the external photons could come from many components including the BLR and have not constant energy density (Bai et al. 2009; Cao & Bai, 2008). (2) The gamma-ray blob is outside the BLR, the external photons from the BLR is Doppler deboosted for the blob. The relation given by equation (4) is invalid, leading no correlation between L_{IC}/L_S and $\nu_{\text{peak}}^{\text{EC}}/\nu_{\text{peak}}^{\text{S}}$, shown in figure 1. We also consider the relation of L_{IC} and L_S , and find an approximate linear relation between them, although the statistical obviousness is not high due to the lack of large sample. This result shows that the ratio of external field and magnetic energy density in the emitting blob is an approximate constant. This implies that the Lorentz factor depends on the external photon field and magnetic field when the blob is inside the BLR. The result indicates an evidence of balancing the jet motion under the action of radiative drag and magnetic derive, and can be used to constrain the jet dynamic model.

Acknowledgements We acknowledge the financial supports from the National Natural Science Foundation of China 10778702, the National Basic Research Program of China (973 Program 2009CB824800), and the Policy Research Program of Chinese Academy of Sciences (KJ CX2-YW-T24).

References

- Abdo, A. A., et al. 2009, *ApJS*, 183, 46
- Abdo A. A., et al. 2010, *ApJ*, 716, 30
- Atwood W. B. et. al. 2009, *ApJ*, 697, 1071
- Bai, J. M., Liu, H. T., & Ma, L. 2009, *ApJ*, 699, 2002
- Begelman M. C., & Rees, M. J. 1984, *MNRAS*, 206, 209
- Blandford, R. D., & Levinson, A. 1995, *ApJ*, 441, 79
- Blażejowski, M., et al. 2000, *ApJ*, 545, 107
- Cao, X., & Bai, J. M. 2008, *ApJ*, 673, L131
- Celotti, A., & Ghisellini, G. 2008, *MNRAS*, 385, 283
- Dermer C. D., Schlickeiser, R., & Mastichiadis, A. 1992, *A&A*, 256, L27
- Dermer C. D., & Schlickeiser, R. 1993, *ApJ*, 416, 458
- Dermer, C. D., Sturnar, S. J., & Schlickeiser, R. 1997, *ApJS*, 109, 103
- Georganopoulos, M., & Kazanas, D. 2003, *ApJ*, 594, L27
- Ghisellini, G., et. al. 1998, *MNRAS*, 301, 451
- Ghisellini, G., & Madau, P. 1996, *MNRAS*, 280, 67
- Ghisellini, G., & Tavecchio, F. 2008, *MNRAS*, 386, L28
- Keppens, R. et al. 2008, *A&A*, 486, 663
- Kubo, H., et al. 1998, *ApJ*, 504, 693
- Leismann, T., et al. 2005, *A&A*, 436, 503
- Liu, H. T., Bai, J. M., & Ma, L. 2008, *ApJ*, 688, 148
- Luo, Q., & Protheroe, R. J. 1999, *MNRAS*, 304, 800
- Maraschi L., Celotti, A., & Ghisellini, G. 1992, *ApJ*, 397, L5
- Marscher A. P., & Gear, W. K. 1985, *ApJ*, 298, 114
- Marscher A. P. 1998, in *IAU Colloq. 164, Radio Emission from Galactic and Extragalactic Compact Sources*, ed. J. M. Wobel. & G. B. Taylor (ASP Conf. Ser. 144; San Francisco: ASP), 25
- McKinney, J. C. 2006, *MNRAS*, 368, 1561
- Padovani P., & Giommi, P. 1995, *ApJ*, 444, 567
- Sambruna R. M., Maraschi, L., & Urry, C. 1996, *ApJ*, 463, 444
- Sikora, M., Begelman, M., & Rees, M. 1994, *ApJ*, 421, 153
- Sikora, M., et. al. 1996, *MNRAS*, 280, 781
- Tavecchio, F., Maraschi, L., & Ghisellini, G. 1998, *ApJ*, 509, 608
- Tavecchio, F., & Ghisellini, G. 2008, *MNRAS*, 386, 945
- Urry, C. M. 1999, *ApJ*, 11, 159

Table 1 List of sources and basic parameters studied in this paper. (1) source LAT names; (2) redshift; (3) synchrotron peak frequency in unit of [Hz]; (4) synchrotron peak flux in unit of [$\text{erg cm}^{-2} \text{s}^{-1}$]; (5) IC peak frequency in unit of [Hz]; (6) IC peak flux in unit of [$\text{erg cm}^{-2} \text{s}^{-1}$]; (7) synchrotron luminosity with [ergs^{-1}]; (8) IC luminosity in unit of [ergs^{-1}]; (9) IC luminosity calculated by the EC model in unit of [ergs^{-1}]; (10) classification.

0FGL (1)	z (2)	$\log \nu_{\text{peak}}^{\text{S}}$ (3)	$\log(\nu F \nu)_{\text{peak}}^{\text{S}}$ (4)	$\log \nu_{\text{peak}}^{\text{IC}}$ (5)	$\log(\nu F \nu)_{\text{peak}}^{\text{IC}}$ (6)	$\log L_{\text{S}}$ (7)	$\log L_{\text{IC}}$ (8)	$\log L_{\text{EC}}$ (9)	Class. (10)
J0137.1+4751	0.859	13.6	-10.8	22.6	-10.6	47.03	46.70	47.29	FSRQ
J0210.8-5100	1.003	12.5	-10.5	22.4	-10.2	47.06	47.73	49.13	FSRQ
J0229.5-3640	2.115	13.5	-10.7	21.8	-10.4	46.73	47.95	45.59	FSRQ
J0238.4+2855	1.213	12.8	-11.7	22.1	-10.8	46.92	47.57	47.79	FSRQ
J0238.6+1636	0.94	13.5	-10.0	23.2	-9.90	47.07	47.50	48.73	BL Lac
J0349.8-2102	2.944	12.9	-10.7	21.8	-10.2	47.35	48.34	47.42	FSRQ
J0423.1-0112	0.916	13.4	-11.3	21.7	-10.3	46.34	47.91	45.20	FSRQ
J0428.7-3755	1.03	13.3	-11.0	22.8	-10.2	46.54	48.14	47.81	BL Lac
J0457.1-2325	1.003	13.1	-10.9	22.8	-9.90	46.88	47.89	48.55	FSRQ
J0531.0+1331	2.07	12.8	-11.0	21.3	-9.80	47.06	48.98	46.33	FSRQ
J0538.8-4403	0.892	13.4	-10.6	22.7	-10.1	47.05	47.22	47.92	BL Lac
J0730.4-1142	1.598	13.1	-11.1	22.6	-10.0	46.98	48.17	48.25	FSRQ
J0855.4+2009	0.306	13.4	-9.8	21.4	-10.5	46.71	46.18	44.97	BL Lac
J0921.2+4437	2.19	13.4	-10.9	22.0	-10.6	47.55	47.73	47.02	FSRQ
J1159.2+2912	0.729	13.1	-11.2	22.0	-10.5	46.58	46.60	46.65	FSRQ
J1229.1+0202	0.158	13.5	-10.7	21.0	-9.60	46.55	46.33	43.82	FSRQ
J1256.1-0548	0.536	12.6	-9.8	22.2	-10.3	45.84	45.74	47.31	FSRQ
J1310.6+3220	0.997	13.1	-10.3	22.5	-10.4	46.99	47.68	48.06	FSRQ
J1457.6-3538	1.424	13.6	-10.9	22.7	-10.2	47.23	48.44	47.69	FSRQ
J1504.4+1030	1.839	13.6	-11.7	22.9	-9.80	45.89	46.70	46.56	FSRQ
J1512.7-0905	0.36	13.1	-10.9	22.3	-9.70	46.56	48.21	47.02	FSRQ
J1522.2+3143	1.487	13.3	-10.6	22.4	-10.2	46.45	47.28	46.71	FSRQ
J1719.3+1746	0.137	13.5	-10.3	24.7	-10.7	44.69	45.84	49.36	BL Lac
J1751.5+0935	0.322	13.1	-10.8	22.2	-10.3	45.95	46.14	46.42	BL Lac
J1849.4+6706	0.657	13.5	-10.6	22.5	-10.5	45.91	45.96	43.98	FSRQ
J2143.2+1741	0.213	14.1	-10.4	22.0	-10.5	47.74	48.23	47.81	FSRQ
J2202.4+4217	0.069	13.6	-10.1	21.9	-10.8	45.28	44.85	44.14	BL Lac
J2254.0+1609	0.859	13.6	-11.5	22.5	-9.80	45.55	46.25	46.21	FSRQ
J2327.3+0947	1.843	13.1	-11.0	21.5	-10.3	47.59	48.24	48.46	FSRQ
J2345.5-1559	0.621	13.3	-9.5	22.5	-10.7	46.85	48.49	45.91	FSRQ
J0033.6-1921	0.61	16.1	-11.1	24.3	-11.1	46.23	46.17	44.90	BL Lac
J0449.7-4348	0.205	15.6	-10.2	23.9	-10.5	46.29	46.17	44.15	BL Lac
J0507.9+6739	0.416	16.6	-10.7	24.3	-10.5	46.35	46.32	44.01	BL Lac
J1015.2+4927	0.2	16.3	-10.5	24.5	-10.6	45.78	46.15	44.44	BL Lac
J1104.5+3811	0.03	16.6	-9.4	25	-9.9	45.16	44.62	44.23	BL Lac
J1653.9+3946	0.033	17.1	-10.3	24.7	-10.5	45.25	44.07	42.71	BL Lac
J2000.2+6506	0.047	16.6	-10	24.7	-10.5	44.57	44.56	43.03	BL Lac
J2158.8-3014	0.116	16	-9.7	23.9	-10.2	46.22	46.00	44.28	BL Lac